

# Is the Mallampati Score Useful for Emergency Department Airway Management or Procedural Sedation?

Steven M. Green, MD\*; Mark G. Roback, MD

\*Corresponding Author. E-mail: [steve@stevegreenmd.com](mailto:steve@stevegreenmd.com), Twitter: [@SteveGreenMD](https://twitter.com/SteveGreenMD).

We review the literature in regard to the accuracy, reliability, and feasibility of the Mallampati score as might be pertinent and applicable to emergency department (ED) airway management and procedural sedation. This 4-level pictorial tool was devised to predict difficult preoperative laryngoscopy and intubation, but is now also widely recommended as a routine screening element before procedural sedation. The literature evidence demonstrates that the Mallampati score is inadequately sensitive for the identification of difficult laryngoscopy, difficult intubation, and difficult bag-valve-mask ventilation, with likelihood ratios indicating a small and clinically insignificant effect on outcome prediction. Although it is important to anticipate that patients may have a difficult airway, there is no specific evidence that the Mallampati score augments or improves the baseline clinical judgment of a standard airway evaluation. It generates numerous false-positive warnings for each correct prediction of a difficult airway. The Mallampati score is not reliably assessed because independent observers commonly grade it differently. It cannot be evaluated in many young children and in patients who cannot cooperate because of their underlying medical condition. The Mallampati score lacks the accuracy, reliability, and feasibility required to supplement a standard airway evaluation before ED airway management or procedural sedation. [Ann Emerg Med. 2019;■:1-9.]

0196-0644/\$-see front matter

Copyright © 2018 by the American College of Emergency Physicians.

<https://doi.org/10.1016/j.annemergmed.2018.12.021>

## INTRODUCTION


The Mallampati score is a graded 4-level pictorial scale (Figure) created to predict difficult intubation before general anesthesia<sup>1-3</sup> and is now routinely used for this purpose in operating rooms worldwide.<sup>4,5</sup> The score has subsequently been extrapolated outside of the operating room to procedural sedation, with a number of specialty societies recommending it as a routine screening element.<sup>6-11</sup> The American Academy of Pediatrics, for example, whose guidelines are “intended for all venues,” including the emergency department (ED), stipulates that the routine pre-sedation health assessment include a Mallampati evaluation.<sup>6</sup> The Mallampati score is widely embedded in electronic medical records (Figure) as a standard assessment step before both general anesthesia and procedural sedation.

This extrapolation of Mallampati scoring to procedural sedation is noteworthy because it applies a tool designed to predict difficult intubation to a different setting in which intubation is never the objective and is essentially never performed. Its proponents assume that the Mallampati score might similarly predict airway and respiratory adverse events during procedural sedation, as well as potential difficulty with bag-valve-mask

ventilation, the key rescue intervention for procedural sedation.

The Mallampati score is intended to supplement, but not replace, the baseline clinical judgment of a general multidimensional airway evaluation.<sup>12-15</sup> This standard airway assessment includes a history and physical inspection of the craniofacial structure to identify risk factors such as short neck, obesity, obstructive sleep apnea, long upper incisors or overbite, restricted mouth opening, micrognathia, macroglossia, laryngomalacia, tonsillar hypertrophy, airway edema, blood or vomit in the airway, cervical immobility, and facial or neck trauma. The quality and reliability of this general airway evaluation will at times be hindered or limited by emergency situations, impaired patient cooperation, and unavailable medical history.

Although meta-analyses of Mallampati score accuracy have been published,<sup>5,16-19</sup> we were unable to identify any clinical review article summarizing the accuracy, feasibility, and reliability of this score in the context of ED airway management or procedural sedation. Accordingly, we performed such a narrative review of the Mallampati score literature as might be pertinent and applicable to the ED setting.



**Figure.** Mallampati score as displayed in the preadmission assessment module of the Epic electronic medical record (Epic, Verona, WI).

## MATERIALS AND METHODS

### Search Strategy

A medical librarian conducted a literature search from January 1985 to October 5, 2018, limited to human subjects and the English language. Our search strategy in PubMed was as follows: “mallampati score”[tiab] OR “mallampati scores”[tiab] OR “mallampati test”[tiab] OR “mallampati class”[tiab] OR “mallampati classification”[tiab] AND (“1985/01/01”[PDAT]: “3000/12/31”[PDAT]) AND “humans”[MeSH Terms] AND English[lang]. Our search strategy in Web of Science was as follows: TOPIC: (“mallampati score” OR “mallampati scores” OR “mallampati test” OR “mallampati class” OR “mallampati classification”); Refined by: LANGUAGES: (ENGLISH); Indexes=SCI-EXPANDED, SSCI, ESCI Timespan=1985-2018. Our search strategy for the Cochrane Database of Systematic Reviews was as follows: (“mallampati score” OR “mallampati scores” OR “mallampati test” OR “mallampati class” OR “mallampati classification”) Filter: from 1985. Our search strategy for EMBASE was as follows: (“mallampati score”/exp OR “mallampati score” OR “mallampati classification”/exp OR “mallampati classification” OR “mallampati test”/exp OR “mallampati test”) AND [english]/lim AND [1985-2018]/py.

We searched references of pertinent articles identified by our search strategy for additional relevant articles. We selected publications with an emphasis on the past 10 years, but we did not exclude commonly referenced and influential older publications.

### HISTORY AND NOMENCLATURE

In 1983, Mallampati described a patient for whom intubation was extremely difficult, and for whom a postanesthesia oral examination revealed that the tongue obscured visualization of the uvula and faucial pillars.<sup>1</sup> He then prospectively assessed 210 preoperative

adults with a 3-level expansion of this evaluation with numeric values—1=normal visualization, 2=obscured uvula but visible faucial pillars, and 3=obscured uvula and faucial pillars—and found that this grading was statistically associated with difficult laryngoscopy.<sup>2</sup>

In 1987, Samssoon and Young<sup>3</sup> expanded this evaluation to the 4-level pictorial classification now in widespread use (Figure), which, when applied after the fact to 13 adults who could not be intubated, was class IV in all except one. In 2001, Ezri et al<sup>20</sup> proposed adding a “class 0”—visualization of the epiglottis—to the score, suggesting it as a reliable predictor of easy laryngoscopy. This modification has not persisted because the primary need driving the scale is the prediction of difficult rather than easy laryngoscopy.

Mallampati,<sup>1</sup> Mallampati et al,<sup>2</sup> and Samssoon and Young<sup>3</sup> performed their evaluations with the patient sitting upright, with the mouth opened widely and tongue protruding. In 1992, Tham et al<sup>21</sup> noted that phonation altered the Mallampati assessment, and it is standard practice to recommend that the patient remain quiet during the evaluation.<sup>20</sup>

Given that the 1985 scale by Mallampati et al<sup>2</sup> had just 3 levels, some authors refer to the Samssoon and Young<sup>3</sup> 4-point adaptation as the “modified Mallampati.”<sup>4,5,22</sup> However, given that the 4-level version has achieved widespread use and acceptance as “the Mallampati score,” for simplicity we will refer to it in that fashion. The Mallampati airway assessment has variously been referred to as a “grade,”<sup>2</sup> “class,”<sup>2,3,20</sup> “classification,”<sup>7,8</sup> “test,”<sup>23</sup> “sign,”<sup>24</sup> “index,”<sup>25</sup> or “score.”<sup>6,9-11</sup> Given that its widespread implementation and coding is as a 1- to 4-point numeric score (Figure), we will refer to it in commonly used fashion as a score with class designations I to IV.

### HOW WELL DOES THE MALLAMPATI SCORE PREDICT DIFFICULT LARYNGOSCOPY AND DIFFICULT TRACHEAL INTUBATION?

Although the Mallampati has 4 possible scores, it has been customary in clinical practice and in research to dichotomize it into classes I and II (low risk) versus classes III and IV (higher risk). This binary risk assessment is then used to predict a research outcome of either difficult laryngoscopy or of difficult intubation. The commonly used standard for difficult laryngoscopy is a Cormack-Lehane direct laryngeal view classification of grade III or IV,<sup>5,23,25-31</sup> although the percentage of glottic opening score has also been used.<sup>32</sup> When the chosen outcome is instead difficult intubation, this has been coded as either a numeric threshold (eg, 3 or more unsuccessful attempts<sup>26</sup>)

or a composite score based on the number of attempts and ultimate intubation failure.<sup>31,33</sup>

### Adults receiving general anesthesia

In the largest single study of adults receiving general anesthesia, Heinrich et al<sup>23</sup> evaluated 102,305 preoperative adults and found that patients with a Mallampati score of III or IV were 6.5 times more likely to have difficult laryngoscopy. Despite this statistical association, however, a higher Mallampati score detected less than half of patients with difficult laryngoscopy, with a sensitivity of 45% and a specificity of 89%. There were 5 false-positive warnings for each correct prediction of a difficult airway.<sup>23</sup>

In the second largest study, Lundstrøm et al<sup>33</sup> evaluated 91,297 adults and found that those with a Mallampati score of III or IV were 3.7 times more likely to have difficult intubation. A higher Mallampati score was 20% sensitive and 94% specific. There were 6 false-positive warnings for each correction prediction of a difficult intubation.

Healy et al<sup>26</sup> performed the third largest Mallampati study, with 80,801 anesthetics, and found that a Mallampati score of III or IV was 43% sensitive and 83% specific in predicting difficult laryngoscopy. There were 14 false-positive warnings for each correct prediction of a difficult airway. The Mallampati score was 32% sensitive and 82% specific for predicting difficult intubation (defined as 3 or more attempts), with 52 false-positive warnings for each correct prediction.

Min et al<sup>32</sup> performed a “gray zone” analysis on 263 preoperative adults, in which gray zone results corresponded to a prediction not precise enough for a diagnostic decision. They noted that 72% of Mallampati assessments overall were in this indeterminate gray zone, as were 63% of the subset with difficult laryngoscopy.

Numerous smaller studies reported similar results.<sup>5,16-19</sup> In a 2018 Cochrane meta-analysis, Roth et al<sup>5</sup> identified 80 studies reporting data on the Mallampati score and difficult laryngoscopy, totaling 232,939 participants with 10,545 cases of difficult laryngoscopy. Among the studies, the accuracy varied widely, with a summary sensitivity of 53% (95% confidence interval [CI] 47% to 59%) and specificity of 80% (95% CI 74% to 85%). Overall, there were 8 false-positive warnings for each correct prediction. In this same review, the authors found 24 studies reporting data on the Mallampati score and difficult intubation, totaling 191,849 participants with 6,615 cases of difficult intubation. Among these studies, the accuracy varied widely, with a summary sensitivity of 51% (95% CI 40% to 61%) and a summary specificity of 87% (95% CI 82% to 91%). There were 7 false-positive warnings for each correct prediction.

Roth et al<sup>5</sup> concluded that “[s]creening tests are expected to have high sensitivities. We found that all investigated index tests [including Mallampati score] had relatively low sensitivities with high variability. In contrast, specificities were consistently and markedly higher than sensitivities across all tests. The standard bedside airway examination tests should be interpreted with caution, as they do not appear to be good screening tests.”

An earlier meta-analysis by Lundstrøm et al<sup>16</sup> evaluated 55 studies with 177,088 patients that reported Mallampati score and difficult intubation. They found a summary sensitivity of 35% (95% CI 34% to 36%) and summary specificity of 91% (95% CI 91% to 91%) and concluded that “the modified Mallampati score is inadequate as a stand-alone test.”

Other earlier systematic reviews and meta-analyses reported similar results.<sup>17-19</sup> An American Society of Anesthesiologists task force concluded in 2013 that “there is insufficient published evidence to evaluate the predictive value of multiple features of the airway physical examination versus single features in predicting the presence of a difficult airway.”<sup>15</sup>

### Children receiving general anesthesia

We were able to identify 6 studies of Mallampati score in preoperative children. In the largest, Heinrich et al<sup>28</sup> evaluated 7,551 preoperative children for whom a Mallampati score was documented. They did not provide sufficient data to calculate sensitivity or specificity, but reported that difficult laryngoscopy was encountered in just 6.4% of patients with a Mallampati score of III to IV (ie, 16 false-positive warnings for each correct prediction).

Inal et al<sup>27</sup> studied 250 children for whom a Mallampati assessment was possible and found that a Mallampati score of III to IV was 77% sensitive and 96% specific in predicting difficult intubation.

Rafique and Khan<sup>29</sup> evaluated 116 children who could comply with the Mallampati assessment and found that a higher Mallampati score predicted 1 of the 7 children with difficult laryngoscopy (sensitivity 14%). Data to calculate specificity were not provided.

After excluding children with an American Society of Anesthesiologists physical status greater than 1 and those with anatomic malformation, genetic syndromes, or cognitive deficit, Santos et al<sup>30</sup> enrolled children aged 4 to 8 years who agreed to participate in a “game” in which they would open the mouth and show the tongue. They noted a Mallampati score sensitivity of 76% and specificity of 96% for predicting difficult laryngoscopy in 108 children.

Aggarwal et al<sup>31</sup> studied 82 children aged 1 to 5 years for whom the Mallampati score could be assessed and found

that it was not statistically associated with difficult laryngoscopy or difficult intubation. They did not provide data to calculate a sensitivity or specificity.

Mansano et al<sup>25</sup> studied 43 children younger than 4 years for whom a Mallampati score could be assessed and noted a sensitivity of 62% and specificity of 52% in predicting difficult laryngoscopy.

To our knowledge, there are no meta-analyses of the Mallampati score in children.

### Adults receiving intensive care unit intubation

De Jong et al<sup>34</sup> described 282 obese adults intubated in an ICU, and Mallampati score was reported for 220. In this subset, a Mallampati score of III to IV was 75% sensitive and 88% specific in predicting the 36 patients with difficult intubation.

### Adults receiving ED intubation

Reed et al<sup>35</sup> studied the 86 of 156 adult ED intubations for which Mallampati score could be assessed, and a class III or IV score was 21% sensitive and 67% specific for identifying difficult intubation. There were 6 false-positive warnings for each correct prediction.

Soyuncu et al<sup>36</sup> prospectively studied 366 ED intubations and were unable to evaluate the Mallampati score because its assessment was possible in only 52 instances (14%).

### Summary

The test characteristics for the Mallampati score in predicting difficult laryngoscopy and difficult intubation in adults vary widely from study to study, with poor aggregate sensitivities from the largest meta-analysis of 53% and 51%, respectively. False-positive warnings were frequent in the studies providing data to calculate them, ranging from 5 to 52 for each correct prediction of a difficult airway. Results for children appear similar.

### HOW WELL DOES THE MALLAMPATI SCORE PREDICT DIFFICULT MASK VENTILATION?

Because intubation is never planned and essentially never occurs with procedural sedation, the presumed value of the Mallampati score in this setting is to instead predict difficult bag-valve-mask ventilation, the key rescue intervention for airway and respiratory adverse events during procedural sedation.

### Adults receiving general anesthesia

In a study of 176,679 preoperative adults receiving general anesthesia, Kheterpal et al<sup>37</sup> evaluated the

composite outcome of concurrent difficult bag-valve-mask ventilation and difficult laryngoscopy, an outcome that occurred in just 0.4% of their sample. A Mallampati score of III or IV was 44% sensitive and 85% specific, with 86 false-positive warnings for each correct prediction. These authors did not separately report results for difficult bag-valve-mask ventilation and difficult laryngoscopy.

Healy et al<sup>26</sup> evaluated 80,801 adults and found that a Mallampati score of III or IV was 31% sensitive and 83% specific in predicting difficult bag-valve-mask ventilation. There were 24 false-positive warnings for each correct prediction.

In a study of 53,041 adults, Kheterpal et al<sup>38</sup> found that a Mallampati score of III to IV was 30% sensitive and 89% specific in predicting the 77 patients for whom bag-valve-mask ventilation was “impossible.” There were 259 false-positive warnings for each correct prediction. The score was less predictive than male sex or a history of sleep apnea.<sup>38</sup> In an earlier study by Kheterpal et al,<sup>39</sup> 350 of 22,660 adults (1.5%) had difficult bag-valve-mask ventilation. A Mallampati score of III to IV was 22% sensitive and 89% specific, with 32 false-positive warnings for each correct prediction.

Leoni et al<sup>40</sup> studied 309 obese preoperative adults and found that difficult bag-valve-mask ventilation was 2.1 times as common when the Mallampati score was III or IV. They did not provide data to calculate sensitivity and specificity.

Other smaller studies also noted a low sensitivity for the Mallampati score.<sup>41-43</sup> In a 2018 Cochrane review (which did not include the study by Healy et al<sup>26</sup> discussed above), Roth et al<sup>5</sup> found 6 studies reporting data on the Mallampati score and difficult bag-valve-mask ventilation, totaling 56,232 participants with 493 cases of difficult ventilation. Among the studies, the sensitivity and specificity varied widely, with a summary sensitivity of 17% (95% CI 6% to 39%) and specificity of 90% (95% CI 81% to 95%). There were 66 false-positive warnings for each correct prediction.

### Children receiving general anesthesia

Aggarwal et al<sup>31</sup> studied 82 preoperative children aged 1 to 5 years who were receiving general anesthesia and for whom the Mallampati score could be assessed, and found that this score was not statistically associated with difficult bag-valve-mask ventilation. They did not report data sufficient to calculate sensitivity or specificity.

### Summary

The test characteristics for the Mallampati score in predicting difficult bag-valve-mask ventilation in adults are extremely poor, with an aggregate sensitivity from the

largest meta-analysis of 17%. False-positive warnings were frequent, ranging from 24 to 259 for each correct prediction of difficult bag-valve-mask ventilation. There are inadequate data to confirm or refute the ability of the Mallampati score to predict difficult bag-valve-mask ventilation for children.

### HOW WELL DOES THE MALLAMPATI SCORE PREDICT ADVERSE EVENTS DURING PROCEDURAL SEDATION?

Because adverse events during procedural sedation are often related to partial airway obstruction, it is also plausible that the Mallampati score might predict these overall complications.

#### Adults

We identified 3 studies pertaining to adults. Khan et al<sup>44</sup> studied 258 obese adults receiving fentanyl-midazolam moderate sedation for bronchoscopy and noted a progressive increase in the frequency of transient hypoxemia with increasing Mallampati score (37%, 45%, 48%, and 62% for the 4 Mallampati scores, respectively). However, they reported no further complications or interference with the procedure.

Hung et al<sup>45</sup> reported an endoscopy and colonoscopy case-control study of 45 adults who received naloxone reversal, flumazenil reversal, or both compared with 90 controls who did not. In a multivariable analysis, they noted that reversal patients were 5.0 times more likely (95% CI 2.1 to 11.7) to have a higher Mallampati score than controls, among multiple other significant predictor variables. Raw Mallampati score data were not provided, and thus it is unclear whether these are comparisons of Mallampati score I versus II or whether patients with scores III or IV were even included.

Agostoni et al<sup>46</sup> reviewed 17,542 gastrointestinal endoscopy procedures facilitated by anesthesiologist-administered propofol (“monitored anesthesia care”) and found that the occurrence of overall adverse events was 1.3 times higher (95% CI 1.0 to 1.7) when the Mallampati score was III or IV. The Mallampati score was not an independent predictor of oxygen desaturation or other more serious complications, however. Data were not provided to permit calculations of sensitivity and specificity.

#### Children

In the only pediatric study, Iyer et al<sup>22</sup> evaluated 458 ED children for whom the Mallampati score could be assessed and found a statistically similar frequency of adverse events between those with higher (III or IV) and lower (I or II) Mallampati scores (24% versus 18%).

### Summary

There are inadequate data to confirm or refute an association between the Mallampati score and procedural sedation-related adverse events.

### HOW RELIABLE IS THE MALLAMPATI SCORE?

An extremely important attribute for any test or clinical evaluation is reliability (ie, that independent assessments by qualified observers are likely to obtain the same result). Studies of such interrater reliability have noted Mallampati score agreement in between 42% and 89% of preoperative<sup>21,24,47-53</sup> or dental clinic<sup>54</sup> adults (Table).  $\kappa$  Statistics fall well short of strong agreement (Table).

Wilson and John<sup>24</sup> had 9 anesthetists independently assess the Mallampati score for 10 colleagues and found that they concurred for only a single individual. There were 2 different ratings for 5 individuals and 3 different ratings for 3 individuals, and a last individual received each of the 4 Mallampati scoring options from at least 1 anesthetist.

Lopez et al<sup>48</sup> evaluated interrater reliability (Table) and also assessed intrarater reliability by having the Mallampati score assessed at different times by the same physician for 20 adults. Clinicians agreed with themselves 65% of the time, with a  $\kappa$  of 0.42.

### Summary

The interrater reliability of the Mallampati score evaluation is moderate at best.

### HOW FEASIBLE IS ROUTINE MALLAMPATI ASSESSMENT?

A cooperative patient is required to open the mouth and protrude the tongue, and so the Mallampati assessment can be impossible for patients who cannot or will not comply.

**Table.** Interobserver agreement in distinguishing Mallampati score I or II versus III or IV.\*

Study	Adult Subjects	Agreement, %	$\kappa$
Menon, 2017 <sup>47</sup>	323	83	0.64
Lopez, 2014 <sup>48</sup>	108	42	0.10
Kandray, 2013 <sup>54</sup>	234	77	0.54
Adamus, 2011 <sup>49</sup>	101	89	0.50
Eberhart, 2005 <sup>50</sup>	1,120	Not reported	0.59
Karkouti, 1996 <sup>51</sup>	59	68	0.31
Tham, 1992 <sup>21</sup>	64	58	Not reported
Oates, 1991 <sup>52</sup>	232	84	Not reported
Wilson, 1990 <sup>24</sup>	10	60	Not reported

\*The Table omits one identified study<sup>39</sup> in which analyses of interobserver agreement are presented for the subsets of patients with known difficult and known easy airways, but omitting an overall analysis.

### Critical illness

Soyuncu et al<sup>36</sup> prospectively studied 366 adult ED intubations and found Mallampati assessment possible in only 14%. In a second ED sample, Levitan et al<sup>13</sup> reviewed 838 intubations and found that 68% of patients could not follow simple commands or were in a cervical spine collar, thus rendering a Mallampati evaluation impossible. The Mallampati score could not be assessed in the 3 patients with intubation failure.

Bair et al<sup>14</sup> described 26% of 296 ED adults being urgently intubated as able to comply with the Mallampati evaluation.<sup>14</sup> Reed et al<sup>35</sup> found that the Mallampati score could be assessed in 55% of 156 adult ED intubations overall and for 44% of the 42 patients with a difficult intubation.

In a prospective study of 1,000 adults intubated in an ICU, the Mallampati score could be first assessed in 77%.<sup>55</sup> In a second sample of 282 obese adults intubated in an ICU, 78% could comply with the Mallampati evaluation.<sup>34</sup> A third critical care study of difficult intubation intentionally did not include Mallampati scoring because of its difficulty of assessment in this population.<sup>56</sup>

### Children

Studies of pediatric anesthesia frequently note an inability to assess the Mallampati score. Mansano et al<sup>25</sup> found that 29% of children younger than 4 years were able to cooperate sufficiently to permit the Mallampati evaluation.<sup>25</sup> Aggarwal et al<sup>31</sup> successfully evaluated the Mallampati score in 82% of 100 children aged 1 to 5 years, with all failures in children younger than 3 years. In a review of 11,219 general anesthetics in children, the documentation rate for the Mallampati score was greater than 80% for children older than 6 years and less than 50% for those younger than 3 years.<sup>28</sup> A prospective study of 196 “cooperative, non-irritable” children undergoing elective surgery found that the Mallampati evaluation was possible in 59%, with most failures in preschool children.<sup>29</sup> In a study of ED procedural sedation in children, the Mallampati score was requested but not documented in 20% of 575 patients, again primarily in the younger portion of their sample.<sup>22</sup>

### Summary

The Mallampati evaluation is frequently impossible to assess in critical illness and for children because of inadequate patient cooperation.

### DISCUSSION

We reviewed the literature in regard to the accuracy, reliability, and feasibility of the Mallampati score, outlining

evidence as might be pertinent and applicable to ED airway management and procedural sedation. In each of these areas, we identified important challenges and limitations to the Mallampati score. Although it is important to anticipate that patients may have a difficult airway, there is no specific evidence that the Mallampati score augments or improves the baseline clinical judgment of a standard airway evaluation (details of which are outlined in the “Introduction”). ED airway management is urgent, and the routine requirement for an added Mallampati assessment might delay care and distract from other airway preparations.

To be clinically valuable, a tool to predict an uncommon outcome such as a difficult airway must be highly sensitive. According to aggregate data from the largest meta-analysis,<sup>5</sup> the Mallampati score is 53% sensitive for difficult laryngoscopy and 17% sensitive for difficult bag-valve-mask ventilation. Thus, reliance on the Mallampati score as a screening tool would miss approximately half of the occurrences of difficult laryngoscopy and most of those of difficult bag-valve-mask ventilation while generating numerous false-positive warnings. According to these same aggregate data,<sup>5</sup> the positive and negative likelihood ratios for difficult laryngoscopy are 2.7 and 0.59, respectively. For difficult bag-valve-mask ventilation, they are 1.7 and 0.92. Each of these likelihood ratios is consistent with a trivial and clinically unimportant effect on outcome probability.

Given this weak test performance, it is surprising that the Mallampati score has found such widespread application within modern anesthesia practice. Indeed, the conclusions of meta-analyses<sup>5,16,18,19</sup> and other reviews<sup>4,15</sup> are consistently negative. Although frequent false-positive warnings may not be viewed as detrimental to safe care, their prevalence further supports our conclusion that when treating individual patients, clinicians must be equally concerned and prepared for a difficult airway regardless of the Mallampati score.

The accuracy calculations above consider the Mallampati score as a stand-alone test, rather than its actual clinical use as an adjunct to the standard general airway evaluation. The purpose of any scoring system or clinical decision rule is to augment and improve clinical judgment, not to replace it.<sup>57-60</sup> We were unable to identify any studies of such actual combined application. Any apparent predictive value of the Mallampati score may have already been captured in the standard general airway assessment.<sup>57-60</sup> There is thus no literature evidence to directly support the contention that Mallampati score augments or improves the standard clinical judgment of the general airway evaluation alone.

There is also no evidence that the Mallampati score—a tool designed for predicting difficult intubation—has any

role for the separate indication of procedural sedation. The key rescue intervention for rare airway and ventilatory adverse events resulting from procedural sedation is bag-valve-mask ventilation, for which, as noted above, the Mallampati score exhibits a weak aggregate positive likelihood ratio of 1.7. The few existing studies of Mallampati score for procedural sedation are inadequate and inconclusive.<sup>22,44-46</sup> In their most recent procedural sedation policy, the American College of Emergency Physicians (ACEP) concluded that “[t]here is no evidence that the addition of the Mallampati score to a standard general evaluation of the anatomic or physiologic variants...has any impact on clinical outcomes during sedation, and thus it cannot be recommended.”<sup>61</sup>

In conclusion, the literature evidence demonstrates that the Mallampati score is inadequately sensitive for the identification of difficult laryngoscopy, difficult intubation, and difficult bag-valve-mask ventilation, with likelihood ratios indicating a small and clinically insignificant effect on outcome prediction. Although it is important to anticipate that patients may have a difficult airway, there is no specific evidence that the Mallampati score augments or improves the baseline clinical judgment of a standard airway evaluation. It generates numerous false-positive warnings for each correct prediction. The score is not reliably assessed because independent observers commonly grade it differently. It cannot be evaluated in many young children and in patients who cannot cooperate because of their underlying medical condition. We conclude that the Mallampati score lacks the accuracy, reliability, and feasibility required to supplement a standard airway evaluation before ED airway management or procedural sedation.

*The authors acknowledge Shanalee Tamares, MLIS, for her expert performance of the literature search.*

**Supervising editor:** Henry E. Wang, MD, MS. Specific detailed information about possible conflict of interest for individual editors is available at <https://www.annemergmed.com/editors>.

**Author affiliations:** From the Department of Emergency Medicine, Loma Linda University Medical Center and Children’s Hospital, Loma Linda, CA (Green); and the Departments of Pediatrics and Emergency Medicine, University of Minnesota, Minneapolis, MN (Roback).

**Author contributions:** SMG and MGR conceived the review, drafted the manuscript, contributed to its many revisions, and approved the final manuscript. SMG takes responsibility for the article as a whole.

All authors attest to meeting the four [ICMJE.org](http://www.icmje.org) authorship criteria: (1) Substantial contributions to the conception or design of the

work; or the acquisition, analysis, or interpretation of data for the work; AND (2) Drafting the work or revising it critically for important intellectual content; AND (3) Final approval of the version to be published; AND (4) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

**Funding and support:** By *Annals* policy, all authors are required to disclose any and all commercial, financial, and other relationships in any way related to the subject of this article as per ICMJE conflict of interest guidelines (see [www.icmje.org](http://www.icmje.org)). The authors have stated that no such relationships exist.

**Publication dates:** Received for publication November 19, 2018. Revisions received November 19, 2018; December 13, 2018, and December 18, 2018. Accepted for publication December 20, 2018.

Dr. Wang was the supervising editor on this article. Dr. Green did not participate in the editorial review or decision to publish this article.

## REFERENCES

- Mallampati SR. Clinical sign to predict difficult tracheal intubation (hypothesis). *Can Anaesth Soc J*. 1983;30:316-317.
- Mallampati SR, Gatt SP, Gugino LD, et al. A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc J*. 1985;32:429-434.
- Samsoon GL, Young JR. Difficult tracheal intubation: a retrospective study. *Anaesthesia*. 1987;42:487-490.
- Baker P. Assessment before airway management. *Anesthesiol Clin*. 2015;33:257-278.
- Roth D, Pace NL, Lee A, et al. Airway physical examination tests for detection of difficult airway management in apparently normal adult patients. *Cochrane Database Syst Rev*. 2018;5:CD008874.
- Cote CJ, Wilson S; American Academy of Pediatrics; American Academy of Pediatric Dentistry; AAP. Guidelines for monitoring and management of pediatric patients before, during, and after sedation for diagnostic and therapeutic procedures: update 2016. *Pediatrics*. 2016;138:e20161212-e20161212.
- Early DS, Lightdale JR, Vargo JJ, et al; ASGE Standards of Practice Committee. Guidelines for sedation and anesthesia in GI endoscopy. *Gastrointest Endosc*. 2018;87:327-337.
- Dumonceau J-M, Riphaut A, Schreiber F, et al. Non-anesthesiologist administration of propofol for gastrointestinal endoscopy: European Society of Gastrointestinal Endoscopy, European Society of Gastroenterology and Endoscopy Nurses and Associates Guideline—updated June 2015. *Endoscopy*. 2015;47:1175-1189.
- Olsen JW, Barger RL, Doshi SK. Moderate sedation: what radiologists need to know. *AJR Am J Roentgenol*. 2013;201:941-946.
- Royal College of Emergency Medicine. Best practice guideline—pharmacological agents for procedural sedation and analgesia in the emergency department. Version 2017. Available at: [https://www.rcem.ac.uk/docs/College%20Guidelines/Pharmacological%20Agents%20for%20Procedural%20Sedation%20and%20Analgesia%20\(Jan%202017%20Revised\).pdf](https://www.rcem.ac.uk/docs/College%20Guidelines/Pharmacological%20Agents%20for%20Procedural%20Sedation%20and%20Analgesia%20(Jan%202017%20Revised).pdf). Accessed October 7, 2018.
- Coté CJ, Wilson S. Guidelines for monitoring and management of pediatric patients before, during, and after sedation for diagnostic and therapeutic procedures: update 2016. *Pediatr Dent*. 2016;38:13-39.
- Sakles JC, Douglas MJK, Hypes CD, et al. Management of patients with predicted difficult airways in an academic emergency department. *J Emerg Med*. 2017;53:163-171.

13. Levitan RM, Everett WW, Ochroch EA. Limitations of difficult airway prediction in patients intubated in the emergency department. *Ann Emerg Med.* 2004;44:307-313.
14. Bair AE, Caravelli R, Tyler K, et al. Feasibility of the preoperative Mallampati airway assessment in emergency department patients. *J Emerg Med.* 2010;38:677-680.
15. Apfelbaum JL, Hagberg CA, Caplan RA, et al. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology.* 2013;118:251-270.
16. Lundström LH, Vester-Andersen M, Møller AM, et al. Danish Anaesthesia Database. Poor prognostic value of the modified Mallampati score: a meta-analysis involving 177 088 patients. *Br J Anaesth.* 2011;107:659-667.
17. Shiga T, Wajima Z, Inoue T, et al. Predicting difficult intubation in apparently normal patients: a meta-analysis of bedside screening test performance. *Anesthesiology.* 2005;103:429-437.
18. Lee A, Fan LT, Gin T, et al. A systematic review (meta-analysis) of the accuracy of the Mallampati tests to predict the difficult airway. *Anesth Analg.* 2006;102:1867-1878.
19. Vannucci A, Cavallone LF. Bedside predictors of difficult intubation: a systematic review. *Minerva Anesthesiol.* 2016;82:69-83.
20. Ezri T, Wartens RD, Szmuk P, et al. The incidence of class "zero" airway and the impact of Mallampati score, age, sex, and body mass index on prediction of laryngoscopy grade. *Anesth Analg.* 2001;93:1073-1075.
21. Tham EJ, Gildersleve CD, Sanders LD, et al. Effects of posture, phonation and observer on Mallampati classification. *Br J Anaesth.* 1992;68:32-38.
22. Iyer MS, Pitetti RD, Vitale M. Higher Mallampati scores are not associated with more adverse events during pediatric procedural sedation and analgesia. *West J Emerg Med.* 2018;19:430-436.
23. Heinrich S, Birkholz T, Irouschek A, et al. Incidences and predictors of difficult laryngoscopy in adult patients undergoing general anesthesia: a single-center analysis of 102,305 cases. *J Anesth.* 2013;27:815-821.
24. Wilson ME, John R. Problems with the Mallampati sign. *Anaesthesia.* 1990;45:486-487.
25. Mansano AM, Módolo NSP, Silva LMD, et al. Bedside tests to predict laryngoscopic difficulty in pediatric patients. *Int J Pediatr Otorhinolaryngol.* 2016;83:63-68.
26. Healy DW, LaHart EJ, Peoples EE, et al. A comparison of the Mallampati evaluation in neutral or extended cervical spine positions: a retrospective observational study of >80 000 patients. *Br J Anaesth.* 2016;116:690-698.
27. Inal MT, Memis D, Sahin SH, et al. Comparison of different tests to determine difficult intubation in pediatric patients. *Rev Bras Anesthesiol.* 2014;64:391-394.
28. Heinrich S, Birkholz T, Ihmsen H, et al. Incidence and predictors of difficult laryngoscopy in 11,219 pediatric anesthesia procedures. *Paediatr Anaesth.* 2012;22:729-736.
29. Rafique NB, Khan FA. Comparison of Mallampati test, thyromental distance and distance from tragus to nares for predicting difficult intubation in pediatric patients. *Open J Anesthesiol.* 2014;4:104-109.
30. Santos AP, Mathias LA, Gozzani JL, et al. Difficult intubation in children: applicability of the Mallampati index. *Rev Bras Anesthesiol.* 2011;61(84-87):156-162.
31. Aggarwal A, Sharma KR, Verma UC. Evaluation of difficult airway predictors in pediatric population as a clinical investigation. *J Anesth Clin Res.* 2012;3:1-5.
32. Min JJ, Kim G, Kim E, et al. The diagnostic validity of clinical airway assessments for predicting difficult laryngoscopy using a grey zone approach. *J Int Med Res.* 2016;44:893-904.
33. Lundström LH, Møller AM, Rosenstock C, et al. High body mass index is a weak predictor for difficult and failed tracheal intubation: a cohort study of 91,332 consecutive patients scheduled for direct laryngoscopy registered in the Danish Anesthesia Database. *Anesthesiology.* 2009;110:266-274.
34. De Jong A, Molinari N, Pouzeratte Y, et al. Difficult intubation in obese patients: incidence, risk factors, and complications in the operating theatre and in intensive care units. *Br J Anaesth.* 2015;114:297-306.
35. Reed MJ, Dunn MJG, McKeown DW. Can an airway assessment score predict difficulty at intubation in the emergency department? *Emerg Med J.* 2005;22:99-102.
36. Soyuncu S, Eken C, Cete Y, et al. Determination of difficult intubation in the ED. *Am J Emerg Med.* 2009;27:905-910.
37. Kheterpal S, Healy D, Aziz MF, et al. Incidence, predictors, and outcome of difficult mask ventilation combined with difficult laryngoscopy: a report from the multicenter perioperative outcomes group. *Anesthesiology.* 2013;119:1360-1369.
38. Kheterpal S, Martin L, Shanks AM, et al. Prediction and outcomes of impossible mask ventilation: a review of 50,000 anesthetics. *Anesthesiology.* 2009;110:891-897.
39. Kheterpal S, Han R, Tremper KK, et al. Incidence and predictors of difficult and impossible mask ventilation. *Anesthesiology.* 2006;105:885-891.
40. Leoni A, Arlati S, Ghisi D, et al. Difficult mask ventilation in obese patients: analysis of predictive factors. *Minerva Anesthesiol.* 2014;80:149-157.
41. Langeron O, Masso E, Huraux C, et al. Prediction of difficult mask ventilation. *Anesthesiology.* 2000;92:1229-1236.
42. El-Ganzouri AR, McCarthy RJ, Tuman KJ, et al. Preoperative airway assessment: predictive value of a multivariate risk index. *Anesth Analg.* 1996;82:1197-1204.
43. Cattano D, Panicucci E, Paolicchi A, et al. Risk factors assessment of the difficult airway: an Italian survey of 1956 patients. *Anesth Analg.* 2004;99:1774-1779.
44. Khan I, Chatterjee AB, Bellinger CR, et al. Sedation for bronchoscopy and complications in obese patients. *Respiration.* 2016;92:158-165.
45. Hung A, Marshall J, Barnett S, et al. Risk factors and outcomes of reversal agent use in moderate sedation during endoscopy and colonoscopy. *J Clin Gastroenterol.* 2016;50:e25-e29.
46. Agostoni M, Fanti L, Gemma M, et al. Adverse events during monitored anesthesia care for GI endoscopy: an 8-year experience. *Gastrointest Endosc.* 2011;74:266-275.
47. Menon SM, Sampangiramaiah S, Mathew M. Cross sectional observational study performed to see for relation of Mallampati score and extended Mallampati score with body mass index. *J Clin Diagn Res.* 2017;11:UG01-UG03.
48. Lopez KT, Theivanayagam S, Asombang AW, et al. Airway assessment of patients undergoing endoscopic procedures. *South Med J.* 2014;107:764-767.
49. Adamus M, Jor O, Vavreckova T, et al. Inter-observer reproducibility of 15 tests used for predicting difficult intubation. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub.* 2011;155:275-281.
50. Eberhart LHJ, Arndt C, Cierpka T, et al. The reliability and validity of the upper lip bite test compared with the Mallampati classification to predict difficult laryngoscopy: an external prospective evaluation. *Anesth Analg.* 2005;101:284-289.
51. Karkouti K, Rose DK, Ferris LE, et al. Inter-observer reliability of ten tests used for predicting difficult tracheal intubation. *Can J Anaesth.* 1996;43:554-559.
52. Oates JD, Macleod AD, Oates PD, et al. Comparison of two methods for predicting difficult intubation. *Br J Anaesth.* 1991;66:305-309.

53. Rosenstock C, Gillesberg I, Gatke MR, et al. Inter-observer agreement of tests used for prediction of difficult laryngoscopy/tracheal intubation. *Acta Anaesthesiol Scand*. 2005;49:1057-1062.
54. Kandray DP, Juruaz D, Yacavone M, et al. Inter-rater reliability of the Mallampati classification for patients in a dental hygiene clinic. *J Dent Hyg*. 2013;87:134-139.
55. De Jong A, Molinari N, Terzi N, et al; AzuRea Network for the Frida-Rea Study Group. Early identification of patients at risk for difficult intubation in the intensive care unit: development and validation of the MACOCHA score in a multicenter cohort study. *Am J Respir Crit Care Med*. 2013;187:832-839.
56. Joshi R, Hypes CD, Greenberg J, et al. Difficult airway characteristics associated with first-attempt failure at intubation using video laryngoscopy in the intensive care unit. *Ann Am Thorac Soc*. 2017;14:368-375.
57. Schriger DL, Elder JW, Cooper RJ. Structured clinical decision aids are seldom compared with subjective physician judgment, and are seldom superior. *Ann Emerg Med*. 2017;70:338-344.e3.
58. Green SM, Schriger DL, Yealy DM. Methodologic standards for interpreting clinical decision rules in emergency medicine: 2014 update. *Ann Emerg Med*. 2014;64:286-291.
59. Schriger DL, Newman DH. Medical decisionmaking: let's not forget the physician. *Ann Emerg Med*. 2012;59:219-220.
60. Green SM, Schriger DL. The sinking STONE: what a failed validation can teach us about clinical decision rules. *Ann Emerg Med*. 2016;67:433-436.
61. Green SM, Roback MG, Krauss BS, et al. Unscheduled procedural sedation: a multidisciplinary consensus practice guideline. *Ann Emerg Med*. In press.