



Dermatome Leg




Dermatome Arm

Glasgow Coma Scale

| Eye Opening (E) | Verbal Response (V) | Motor Response (M) |
|--|--|--|
| 4 = Spontaneous 3 = To voice 2 = To pain 1 = None | 5 = Normal conversation 4 = Disoriented conversation 3 = Words, but not coherent 2 = No words . . . only sounds 1 = None | 6 = Normal 5 = Localizes to pain 4 = Withdraws to pain 3 = Decorticate posture 2 = Decerebrate 1 = None |
| Total = E + V + M | | |

EKG Territory Chart

| | | | |
|--------------|--------------|-------------|-------------|
| I Lateral | aVR | VI Septal | V4 Anterior |
| II Inferior | aVL Lateral | V2 Septal | V5 Lateral |
| III Inferior | aVF Inferior | V3 Anterior | V6 Lateral |



INTEGRIS
Southwest Medical Center

Emergency Medicine Residency
<http://integrisok.com/emergency-residency>
405-636-7195

Peds Quick Reference

Intubation Dosages

Etomidate 0.3 mg/kg

Ketamine 2 mg/kg

Propofol 1.5-3 mg/kg

Succinylcholine 1.5-2 mg/kg

Rocuronium 1 mg/kg

Resuscitation Dosages

Epinephrine (1:10,000) 0.01 mg/kg Max of 1mg

Bicarbonate 1 mEq/kg

Cardioversion(synchronized) 1st dose 0.5-1 J/kg

2nd dose 2 J/kg

Defibrillation 1st dose 2 J/kg

2nd dose 4 J/kg

Subsequent doses 4-10 J/kg

Normal Pediatric Vital Signs by Age

| Age | Heart Rate (bpm) | Respiratory Rate (bpm) | Systolic Blood Pressure (mm Hg) | Diastolic Blood Pressure (mm Hg) |
|---------------|------------------|------------------------|---------------------------------|----------------------------------|
| Newborn | 90 - 180 | 30 - 50 | 60 ± 10 | 37 ± 10 |
| 1 - 5 months | 100 - 180 | 30 - 40 | 80 ± 10 | 45 ± 15 |
| 6 - 11 months | 100 - 150 | 25 - 35 | 90 ± 30 | 60 ± 10 |
| 1 year | 100 - 150 | 20 - 30 | 95 ± 30 | 65 ± 25 |
| 2 - 3 years | 65 - 150 | 15 - 25 | 100 ± 25 | 65 ± 25 |
| 4 - 5 years | 65 - 140 | 15 - 25 | 100 ± 20 | 65 ± 15 |
| 6 - 9 years | 65 - 120 | 12 - 20 | 100 ± 20 | 65 ± 15 |
| 10 - 12 years | 65 - 120 | 12 - 20 | 110 ± 20 | 70 ± 15 |
| 13+ years | 55 - 110 | 12 - 18 | 120 ± 20 | 75 ± 15 |

Anti-Epileptic Drugs

Lorazepam 0.1 mg/kg (may use twice)

Fosphenytoin 20 mg/kg

Levetiracetam 20 mg/kg

Valproic Acid 20 mg/kg

Phenobarbital 20 mg/kg

Propofol 1-2 mg/kg (infuse at 10mg/min)

INTEGRIS

Southwest Medical Center